

AFOSR 820

IMM-NYU 282  
JUNE 1961



NEW YORK UNIVERSITY  
INSTITUTE OF  
MATHEMATICAL SCIENCES

# Extensions and Corollaries of Recent Work on Hilbert's Tenth Problem

MARTIN DAVIS

PREPARED UNDER  
CONTRACT NO. AF49(638)-777  
MATHEMATICAL SCIENCES DIRECTORATE  
AIR FORCE OFFICE OF SCIENTIFIC RESEARCH

REPRODUCTION IN WHOLE OR IN PART  
IS PERMITTED FOR ANY PURPOSE  
OF THE UNITED STATES GOVERNMENT.

IMM-282  
c.1



New York University  
Institute of Mathematical Sciences

EXTENSIONS AND COROLLARIES OF RECENT WORK  
ON HILBERT'S TENTH PROBLEM

Martin Davis

ABSTRACT: The theorem that every recursively enumerable set is exponential Diophantine is improved; a sharp form is given of Kleene's normal form theorem, a problem of Quine is proved recursively unsolvable.

"Qualified requestors may obtain copies of this report from the ASTIA Document Service Center, Arlington Hall Station, Arlington 12, Virginia. Department of Defense contractors must be established for ASTIA services, or have their "need-to-know" certified by the cognizant military agency of their project or contract".

The research reported in this document has been sponsored by the Mathematical Sciences Directorate, Air Force Office of Scientific Research, Washington 25, D. C., under Contract No. AF 49(638)-777.



This paper consists of three separate notes related only in that each of the three either extends or employs the results of [2], with which acquaintance is assumed,

1. A sharpening of Kleene's normal form theorem.

By a form of Kleene's normal form theorem<sup>1</sup> we may understand an assertion stating:

Theorem. There is a function  $U(y)$  and a predicate  $T(z,x,y)$  both belonging to the class  $Q$  such that a function  $f(x)$  is partially computable if and only if for some number  $e$ :

$$f(x) = U(\min_y T(e,x,y)) \quad .$$

In its original form, this result was stated with  $Q$  the class of primitive recursive functions and predicates. It is well-known<sup>2</sup> that smaller classes  $Q$  suffice. We wish to point out here that (assuming variables to range over the positive integers) we may take for  $Q$  the following extremely modest class:

(1) A function  $f$  belongs to  $Q$  if and only if  $f$  can be obtained by repeated application of the operation of composition to the functions:  $2^x$ ,  $x \cdot y$ ,  $N(x) = 0$ ,  $U_i^n(x_1, \dots, x_n) = x_i$ ,  $K(x)$ ,  $L(x)$ , where  $K(x)$ ,  $L(x)$  are recursive pairing functions.

(2) A predicate  $R(x_1, \dots, x_n)$  belongs to  $Q$  if:

$$R(x_1, \dots, x_n) \longleftrightarrow f(x_1, \dots, x_n) = g(x_1, \dots, x_n)$$

where  $f, g \in Q$  .

In fact, we may even take  $U(y) = K(y)$

To see this we begin by noting that by Corollary 5 of [2], (or rather the immediate extension thereof to predicates), we



have:

$$\begin{aligned} \bigvee_y T_2(z, x, u, y) &\longleftrightarrow \bigvee_{x_1, \dots, x_n} P(z, x, u, x_1, \dots, x_n, 2^{x_1}, \dots, 2^{x_n}) = 0 \quad . \\ &\longleftrightarrow \bigvee_{x_1, \dots, x_n} \left\{ \sum_{j=1}^m f_j(z, x, u, x_1, \dots, x_n, 2^{x_1}, \dots, 2^{x_n}) \right. \\ &\quad \left. = \sum_{j=1}^m g_j(z, x, u, x_1, \dots, x_n, 2^{x_1}, \dots, 2^{x_n}) \right\} \end{aligned}$$

where  $f_j, g_j \in Q$ ,  $j = 1, 2, \dots, m$  .

Now, using the fact

$$\begin{aligned} \sum A_j = \sum B_j &\longleftrightarrow \sum 2^{A_j} = \sum 2^{B_j} \quad , \\ &\longleftrightarrow \prod 2^{A_j} = \prod 2^{B_j} \quad , \end{aligned}$$

we see that

$$\bigvee_y T_2(z, x, u, y) \longleftrightarrow \bigvee_{x_1, \dots, x_n} R(z, x, u, y, x_1, \dots, x_n)$$

where  $R \in Q$  .

Now, let  $q_1(t) = K^{n-1}(t)$ ,

$$q_j(t) = L(K^{n-j}(t)) \quad , \quad j = 2, 3, \dots, n \quad ,$$

where the exponent on  $K$  indicates iterated application, so that

$q_j(t) \in Q$ ,  $j = 1, 2, \dots, n$  . Thus

$$\begin{aligned} \bigvee_y T_2(z, x, u, y) &\longleftrightarrow \bigvee_t R(z, x, u, y, q_1(t), \dots, q_n(t)) \\ &\longleftrightarrow \bigvee_t S(z, x, u, y, t) \end{aligned}$$

where  $S \in Q$  .





Let  $f(x)$  be any partially computable function. Then the predicate  $u = f(x)$  is semicomputable (recursively enumerable).

Hence, for some  $e$ ,  $u = f(x) \iff \bigvee_y T_2(e, x, u, y)$

$$\iff \bigvee_t S(e, x, u, t) \quad .$$

Finally,

$$f(x) = K\left(\min_y S(e, x, K(y), L(y))\right) \quad .$$

So, we have derived Kleene's normal form theorem with

$$T(z, x, y) \iff S(z, x, K(y), L(y))$$

and

$$u(y) = K(y) \quad .$$

2. Negative solution to a problem of Quine. In [4], Quine proposed the following problem:

Let us consider schemata constructed from the following ingredients:

Numerals, variables ranging over the non-negative integers, the symbols of sum, product and power,  $=$ , and the truth-function signs.

Such a schema is called valid if it becomes a true sentence whenever all of the variables occurring in it are replaced by numerals. The proposed problem is to give an algorithm for determining whether or not a given schema of this kind is valid.

We note here that the recursive unsolvability of this problem



follows directly from the results of [2]. For, to each exponential Diophantine equation,  $E = F$ , there corresponds, mechanically, a "translation":  $\Gamma = \Delta$  which is a schema of the kind being considered. Moreover,  $E = F$  has a solution if and only if the schema  $\sim(\Gamma = \Delta)$  is not valid. Hence, an algorithm for solving Quine's problem could be used to solve the decision problem for exponential Diophantine equations. But, by [2], there is no algorithm for solving this latter problem. Hence, Quine's problem is likewise unsolvable.

3. Diophantine representation of recursively enumerable sets in terms of a single predicate of exponential growth. A predicate  $\rho(u,v)$  will be called a Julia Robinson predicate if:

- (1)  $\rho(u,v) \rightarrow v \leq u^u$
- (2) For each  $k > 0$ , there are  $u,v$  such that:  

$$\rho(u,v) \wedge v > u^k .$$

We shall prove the:

Theorem. Let  $S$  be a recursively enumerable set. Then, there is a polynomial  $P$  such that:

$$S = \left\{ x \mid \bigvee_{x_1, \dots, x_n, u, v} [P(x, x_1, \dots, x_n, u, v) = 0 \wedge \rho(u, v)] \right\}$$

for every Julia Robinson predicate  $\rho(u,v)$ .

Since, e.g. the predicate  $v = 2^u \wedge u > 1$  is a Julia Robinson predicate, we have the

Corollary 1. Let  $S$  be a recursively enumerable set. Then, for some polynomial  $P$ ,



$$S = \left\{ x \mid \bigvee_{x_1, \dots, x_n, u} P(x, x_1, \dots, x_n, u, 2^u) = 0 \right\} .$$

This generalizes Corollary 5 of [2]. Moreover, the proof of Corollary 6 of [2], if applied to the present Corollary 1 instead of to Corollary 5 of [2], yields:

Corollary 2. For every recursively enumerable set S there is a function  $P(x_1, \dots, x_n, u, 2^u)$ , where P is a polynomial, whose range (for positive integer value of the variables) consists of the members of S together with the non-positive integers.

If in particular we choose for S, the set of positive primes, we obtain a curious "prime-representing" function.

It remains to prove the theorem stated above. In doing so we generalize the methods, relating to Pell's equation, of [5].<sup>3</sup> We recall the notation  $x = a_n$ ,  $y = a'_n$  for the successive solutions of the Pell equation  $x^2 - (a^2 - 1)y^2 = 1$ .

Lemma 1. There is a Diophantine predicate  $\psi(a, u)$  such that:

- (1)  $\psi(a, u) \rightarrow u \geq a^a$
- (2)  $a > 1 \rightarrow \bigvee_u \psi(a, u) .$

Proof. This is a weakening of Lemma 8 of [5].

Lemma 2. There is a Diophantine predicate  $D(c, y, z)$  such that

- (1)  $a > c \wedge D(c, y, z) \rightarrow a > y^z$
- (2)  $\bigwedge_{y, z} \bigvee_c D(c, y, z)$



Proof. Let

$$D(c, y, z) \longleftrightarrow \bigvee_b [b > y \wedge b > z \wedge \psi(b, c)] .$$

Then,

$$a > c \wedge D(c, y, z) \longrightarrow \bigvee_b [a > c \geq b^b > y^z] .$$

Lemma 3. If  $y > 1$  and  $a > y^z$  then  $y^z = [u/a_z]$  where<sup>4</sup>  $u$  is chosen as a solution of

$$u^2 - (a^2 y^2 - 1)v^2 = 1 \text{ for which } a_z \leq u \leq a \cdot a_z .$$

Proof. By Lemma 9 of [5],  $y^z = [(ay)_z/a_z]$ , and by Lemma 10, the number  $u$  is precisely  $(ay)_z$ .

Lemma 4,

$$\bigwedge_{i \leq m} (x_i = y_i^{z_i}) \longleftrightarrow \bigvee_{r_1, \dots, r_m} \bigwedge_{i \leq m} [E(r_i, x_i, y_i, z_i, a) \wedge \bigwedge_{i \leq m} (r_i = a_{z_i})]$$

where  $E$  is a Diophantine predicate and where  $a > c_1, c_2, \dots, c_m, z_1, \dots, z_m$  with the  $c_1, \dots, c_m$  satisfying  $D(c_i, y_i, z_i)$ .

Proof. We need only take

$$E(r_i, x_i, y_i, z_i, a) \longleftrightarrow \bigvee_{u, v} [(u^2 - (a^2 y_i^2 - 1)v^2 = 1)$$

$$\wedge r_i \leq u \leq a \cdot r_i \wedge r_i x_i \leq u < r_i (x_i + 1)] \bigvee [x_i = y_i = 1] .$$

Lemma 5. If  $1 < r < a_a$  and  $a > z$ , then

$$r = a_z \longleftrightarrow \bigvee_s [r^2 - (a^2 - 1)(z + s(a - 1))^2 = 1] .$$





Proof. This follows from Lemma 7 of [5].

Lemma 6.

$$\bigwedge_{i \leq m} (x_i = y_i^{z_i}) \longleftrightarrow \bigvee_{a, d} [F(x_1, \dots, x_m, y_1, \dots, y_m, z_1, \dots, z_m, a, d) \wedge \rho(a, d)]$$

where  $F$  is a Diophantine predicate and  $\rho$  may be any Julia Robinson predicate.

Proof. We claim that, using the notation of Lemma 4:

$$\begin{aligned} \bigwedge_{i \leq m} (x_i = y_i^{z_i}) &\longleftrightarrow \bigvee_{r_1, \dots, r_m} \bigvee_a \left\{ \bigwedge_{i \leq m} [E(r_i, x_i, y_i, z_i, a) \right. \\ &\quad \wedge (a > z_i) \wedge \bigvee_{s_i} [r_i^2 - (a^2 - 1)(z_i + s_i(a - 1))^2 = 1] \\ &\quad \wedge \bigvee_{c_1, \dots, c_m} \left[ \bigwedge_{i \leq m} (D(c_i, y_i, z_i) \wedge a > c_i) \right] \\ &\quad \left. \wedge \bigvee_d [r_1, \dots, r_m \leq d \wedge \rho(a, d)] \right\}. \end{aligned}$$

For, if the right-hand side holds, then  $r_1, \dots, r_m \leq d \leq a^a < a_{a_z}$  so that by Lemma 5,  $r_i = a_{z_i}$ , and finally, by Lemma 6,  $x_i = y_i^{z_i}$ . Conversely, if the left-hand side holds choose  $c_i$  so that  $D(c_i, y_i, z_i)$  is satisfied, then let  $z = \max_{i \leq m} z_i$ , and choose  $a, d$  so

that  $a > c_i$ ,  $a > z$ ,  $\rho(a, d)$ , and  $d > a_z$ . Then,

$$r_i = a_{z_i} \leq a_z < d,$$

and the result follows by Lemma 4 and 5.



Lemma 7. Let  $S$  be a recursively enumerable set. Then, there is a polynomial  $P$  such that:

$$S = \left\{ x \mid \bigvee_{x_1, \dots, x_m} \bigvee_{y_1, \dots, y_m} \bigvee_{z_1, \dots, z_m} [P(x, x_1, \dots, x_n, y_1, \dots, y_m, z_1, \dots, z_m) = 0] \wedge \bigwedge_{i \leq m} (x_i = y_i^{z_i}) \right\} .$$

Proof. This lemma is essentially a restatement of the main result of [2], namely that every recursively enumerable set is exponential Diophantine.

The theorem now follows at once from Lemmas 6 and 7.



Footnotes

1. Cf. [1] or [3].
2. Cf. [3] and [6].
3. However, we are following [2] rather than [5] in taking variables to have the positive integers (rather than the non-negative integers) as their range.
4. [...] here means, as usual, "the greatest integer  $\leq \dots$ ."



References

- [1] Martin Davis, Computability and unsolvability, McGraw-Hill, 1958.
- [2] Martin Davis, Hilary Putnam, and Julia Robinson, The decision problem for exponential Diophantine equations. Annals of Mathematics, forthcoming.
- [3] S. C. Kleene, Introduction to metamathematics, D. Van Nostrand Company, Inc., 1952.
- [4] W. V Quine, On decidability and completeness, Synthese, vol. VII (1948-9), pp. 441-46.
- [5] Julia Robinson, Existential definability in arithmetic, Transactions of the American Mathematical Society, vol. 72 (1952), pp. 437-449.
- [6] Raymond Smullyan, Theory of formal systems, Annals of Mathematics Studies, 1961.





DISTRIBUTION LIST AIR FORCE OFFICE OF SCIENTIFIC RESEARCH  
MATHEMATICAL SCIENCES DIRECTORATE  
(ONE COPY UNLESS OTHERWISE NOTED)

ALABAMA

Commander  
Army Rocket & Guided Missile Agency  
ATTN: ORDXR-OTL  
Redstone Arsenal, Alabama

BELGIUM

Commander (3)  
European Office, ARDC  
47 Rue Cantersteen  
Brussels, Belgium

CALIFORNIA

Applied Mathematics & Statistics  
Laboratory  
Stanford University  
Stanford, California

Department of Mathematics  
University of California  
Berkeley, California

Commander  
Air Force Flight Test Center  
Attn: Technical Library  
Edwards Air Force Base, California

The Rand Corporation (2)  
Technical Library  
1700 Main Street  
Santa Monica, California

Commander  
1st Missile Division  
ATTN: Operations Analysis Office  
Vandenburg Air Force Base,  
California

CONNECTICUT

Department of Mathematics  
Yale University  
New Haven, Connecticut

DISTRICT OF COLUMBIA

Office of Naval Research (2)  
Department of the Navy  
ATTN: Code 432  
Washington 25, D. C.

Director  
Department of Commerce  
Office of Technical Services  
Washington 25, D. C.

Administrator (6)  
National Aeronautics & Space  
Administration  
ATTN: Documents Library  
1520 H Street, N. W.  
Washington 25, D. C.

Library  
National Bureau of Standards  
Washington 25, D. C.

Data Processing Systems Division  
National Bureau of Standards  
ATTN: Mr. Russell A. Kirsch  
Washington 25, D. C.

Applied Mathematics Division  
National Bureau of Standards  
Washington 25, D. C.

Headquarters, USAF  
Assistant for Operations Analysis  
Deputy Chief of Staff,  
Operations, AF00A  
Washington 25, D. C.

Commander (2)  
Air Force Office of Scientific  
Research  
ATTN: SRM  
Washington 25, D. C.



NEBRASKA

Commander  
Strategic Air Command  
ATTN: Operations Analysis  
Offutt Air Force Base  
Omaha, Nebraska

NEW JERSEY

The James Forrestal Research  
Center Library  
Princeton University  
Princeton, New Jersey

Library  
Institute for Advanced Study  
Princeton, New Jersey

Department of Mathematics  
Fine Hall  
Princeton University  
Princeton, New Jersey

Commanding General  
Signal Corps Engineering Laboratory  
ATTN: SIGFM/EL-RPO  
Ft. Monmouth, New Jersey

NEW MEXICO

Commander  
Air Force Missile Development  
Center  
ATTN: Technical Library, HDOI  
Holloman Air Force Base, New Mexico

Commander  
Air Force Special Weapons Center  
ATTN: Technical Library, SWOI  
Kirtland Air Force Base  
Albuquerque, New Mexico

NEW YORK

Professor J. Wolfowitz  
Mathematics Department  
White Hall  
Cornell University  
Ithaca, New York

Department of Mathematics  
Syracuse University  
Syracuse, New York

Institute for Mathematical  
Sciences  
New York University  
ATTN: Professor M. Kline  
25 Waverly Place  
New York 3, New York

Institute for Aeronautical  
Sciences  
ATTN: Librarian  
2 East 64th Street  
New York 16, New York

NORTH CAROLINA

Department of Mathematics  
University of North Carolina  
Chapel Hill, North Carolina

Department of Statistics  
University of North Carolina  
Chapel Hill, North Carolina

Office of Ordnance Research (2)  
Box CM  
Duke Station  
Durham, North Carolina

Department of Mathematics  
Duke University  
Duke Station  
Durham, North Carolina

OHIO

P. O. Box AA  
Wright-Patterson Air Force Base  
Ohio

Commander  
Wright Air Development Division  
ATTN: WCOSI  
Wright-Patterson Air Force Base  
Ohio

Commander  
Aeronautical Research  
Laboratories  
ATTN: Technical Library  
Wright-Patterson Air Force Base  
Ohio



USAF Institute of Technology (2)  
Library  
ATTN: MCLI-ITLIB  
Building 125, Area B  
Wright-Patterson Air Force Base  
Ohio

Mathematics Research Center,  
U. S. Army  
ATTN: R. E. Langer  
University of Wisconsin  
Madison, Wisconsin

#### PENNSYLVANIA

Department of Mathematics  
Carnegie Institute of Technology  
Pittsburgh, Pennsylvania

Department of Mathematics  
University of Pennsylvania  
Philadelphia, Pennsylvania

#### TENNESSEE

AEDC Library  
ARO, Inc.  
Arnold AF Station, Tennessee

U. S. Atomic Energy Commission  
Technical Information Service  
Extension  
P. O. Box 62  
Oak Ridge, Tennessee

#### TEXAS

Applied Mechanics Reviews (2)  
Southwest Research Institute  
8500 Culebra Road  
San Antonio 6, Texas

Department of Mathematics  
Rice Institute  
Houston, Texas

#### VIRGINIA

Armed Services Technical (2)  
Information Agency  
ATTN: TIPDR  
Arlington Hall Station  
Arlington 12, Virginia

#### WISCONSIN

Department of Mathematics  
University of Wisconsin  
Madison, Wisconsin

[illegible]

PRINTED IN U. S. A.



NYU  
IMM-282

c.1

Davis

NYU  
IMM-282

c.1

Davis

NYU  
IMM-282

c.1

Davis

AUTHOR

Extensions and corollaries of

TITLE recent work on  
Hilbert's tenth problem

DATE DUE

BORROWER'S NAME

1972 1976

*M. Spatling*

N. Y. U. Institute of  
Mathematical Sciences

~~25 Waverly Place~~

New York 3, N. Y.

4 Washington Place

